# Determination of Public Land (Rangeland) Health for 64079 WEST SIDE

The Record of Decision (ROD) for the New Mexico Standards for Public Land Health and Guidelines for Livestock Grazing Management (dated January 2001) adopted three Standards for Public Land Health. These are (1) Upland Sites Standard, (2) Biotic Communities, Including Native, Threatened, Endangered, and Special Status Species Standard and (3) Riparian Sites Standard.

The ROD also established a process for the BLM Field Offices for the implementation. Through a public participation process, the Roswell Field Office developed and adopted indicators to use in conjunction with existing monitoring data to assess these standards.

Field assessment worksheets and other available data that evaluate the local indicators were completed for this allotment. Based on the assessments, it is my determination that the public land within the West Side allotment #64079 meets the Upland Sites standard and (2) Biotic Communities, including Native, Threatened, Endangered, and Special Status Species standard. There are no public land Riparian areas on this allotment, therefore this standard will not be addressed.

/s/ T. R. KREAGER
Assistant Field Manager

09/20/2004

er Date

## Standards of Public Land Health Evaluation of 64079 WEST SIDE Allotment [ 07/27/2004 ]

The Roswell Field Office conducted rangeland health assessments at two (2) study sites within the West Side allotment #64079. The assessments looked at the Soil/Site Stability, Hydrologic Function and Biotic Integrity indicators within the vicinity of each study site. Existing monitoring data was incorporated into and in support of the field assessment. The summary of each assessment is attached and shown in the following table.

Study Area	UPLAND			BIOTIC			RIPARIAN		
or Assessment Area	Meets	Monitor an Indicator	Does Not Meet	Meets	Monitor an Indicator	Does Not Meet	Meets	Monitor an Indicator	Does Not Meet
64079- NORTH-F060	X			X			N/A		
64079- SOUTH-F061	X			X			N/A		

Twenty-two (22) indicators for Rangeland Health were evaluated for the public land on the West Side allotment #64079. Ten (10) of these assessed soil site stability, 11 hydrologic function and 13 biotic integrity. These qualitative assessments in conjuntion with quantitaive information gathered from previous data collected on 2 range trend study locations within the allotment were utilized to make rangeland health determinations. Quantitative evaluations are performed by the Roswell Field Office include some or all of the following: ground and vegeative cover and composition, production, frequency, occurence and ecological condition. These collections which were initiated in the early 1980's are scheduled and conducted approximately every 5 years.

North Pasture is a Shallow SD-3 ecological site with 3,242 acres/1474 hectares. The soil phase is a (Tg) Tencee-Upton complex on nearly level to gently rolling soil on upland ridges west of the Pecos River. Slopes are 0 to 9 percent. This allotment possesses habitat for pronghorn (Antilocapra americana) and mule deer (Odocoileus hemionus). At the bottomland portion of this pasture next to and including the private land holding, is where livestock are currently grazing and appear to be holding here. With the recent rainfall, the bottomland swale comprised mainly of alkali sacaton (Sporobolus airoides) and tobosa (Pleuraphis mutica) is vegetating with some puddling and ponding occuring. The study area is however situated upland from the swale and is made up of a desert shrub grassland mix.

The majority of indicators assessed rated Slight to Moderate. Only minor erosion is taking place with stable and short water flow patterns. Approximately 16 percent rock cover has been the long-term average with the remainder made up of 50% bareground

and approximately 18 percent vegetative cover. Indicators with all soil and hydrologic attributes rated Slight to Moderate except for physical crusts which were Moderate. Physical crust is weak in some places and it's continuity is broken. The site has some gravelly influence with small rock and pebbles on the soil surface. The plant groups are holding steady with influxes of Christmas cholla (Opuntia imbricata) and creosote (Larrea tridentata). The black grama (Bouteloua eriopoda) is somewhat abundant with occurences of burrograss (Scleropogon brevifolius), feather dalea (Dalea formosa), range ratany (Krameria spp.) and dogweed (Dyssodia spp.). Annual production is Moderate with 1/2 of the long-term average estimated. Invasive plants rates Moderate also with creosote scattered throughout.

South Pasture rated most indicators in the Slight to Moderate category. The recent dry conditions have possibly had an impact on this site. Bare patches of soil can be observed with some gravel and pebble cover. Functional'structural groups rate Moderate as dropseed (Sporobolus spp.), sideoats grama (Bouteloua curtipendula) and skunkbush (Rhus spp.) have been reduced. Christmas cholla is growing in approximately 2/3 of the treated creosote which suggests that the herbididal treatment is the recent past has helped to propagate this plant. Mesquite (Prosopis glandulosa) can be seen surrounding a dirt tank which is holding water, downhill from this upland area. A gas pipeline is intersecting the site and is evidenced by the rocky berm which has been out into place. Very little vegetation is observed on the pipeline. Annual production rates Moderate as 40% of the ESD average and long-term data is estimated. Invasive plants is Moderate with creoste scattered and snakeweed (Gutierrezia sarothrae) common throughout. All other indicators rated None to Slight with normal ranges of variability.

Hydroogy - Pasture North - The physical/biological crust indicator rated as moderate. The soil crusts were only found in protected areas with a minor component in interspaces. All other indicators rated as none to slight or slight to moderate. Sand and gravel deposits of Quaternary pediment deposits outcrop in the area.

Pasture South - The rills, water flow patterns, pedestals and/or terracettes, bareground, gullies, wind scoured, blowouts, and or deposition areas, litter movement, soil surface resistance to erosion, soil surface loss or degradation, plant community composition and distribution relative to infiltration and runoff, compaction layer, litter amount, and physical/chemical/biological crusts indicators have rated as none to slight or slight to moderate. Sand and gravel deposits of Quaternary alluvial deposits outcrop in the area.

Wildlife - Evaluation of the integrity of the biotic community considered several indicators as attribute indices for the area of interest. Biotic indicators are interrelated with several other indicators, including soil/site stability, hydrologic function, and vegetation. Several indicators are singularly biotic and address the vegetative aspect of the ecological site description, such as functional/structural groups and annual production as discussed above.

North Pasture - Specifically, two biotic indicators fell within the Moderate rating, annual production and invasive plants (creosote being cited). Considering present climate

regimes, annual can be expected to fall within the normal range of variability. Range condition (based on production) appears to have declined over the years, even when considering the drought conditions that has prevailed over the past several years. Browse species can still be found in the drainages that characterize the Walnut Creek and Felix River watershed. As the area of interest falls within an ecotone between the Chihuahuan desert and grasslands biome, desert shrub components can be expected in the area and has increased with declining range site conditions and overall drying conditions over time, especially on the shallow soils.

In addition to the standard worksheet biotic factors, four specific wildlife indicators and descriptors are included in this evaluation. Wildlife habitat and population indicators rate Slight to Moderate, primarily for pronghorn (Antilocapra americana) and a variety of non-game terrestrial species. The composition of vegetation reflects current climatic conditions, e.g., drought for the past several years. Range site production and cover of a variety of preferred plant species for wildlife, such as forbs and woody browse species, and the availability of seed for food and regeneration, is moderated by climate and land use. It should be noted that the study site is on the shallower soils although swales are another important habitat component not specifically addressed in this assessment.

With respect to Special Status Species, none are known to occur in the area of interest at this time and the habitat and population indicators are, therefore, rated None to Slight.

South Pasture - Basically the same as North Pasture with the addition of functional /structural groups in the Moderate rating due to dropseed, sideoats grama and skunkbush falling out of the vegetative composition. Further note that this pasture was chemically treated for creosote some time in the past.

It is the professional opinion of the Assessment Team that the public land within the West Side allotment meets the Upland and Biotic standards. There are no Riparian issues present, therefore this standard was not addressed. See site notes and recommendations for further information regarding these assessments.

**Recommendations:** Regular scheduled monitoring should continue on the two sites. A more critical look at the brush control should be performed to determine the effectiveness of the herbicidal treatment on creosote. It appears that the treatment helped to establish a new crop of Christmas cholla growing up from the middle of the dead creosote clumps. Perhaps a more condusive environment was established for the cholla. A closer observation to record the encroachment of cholla should be done.

RFOs	Upland	and Biotic Standa	rd Asse	ssment S	ummary `	Worksh	eet
		SITE 64079	-NORT	H-F060			
Legal La	and Desc	NWNE 7 0150S 024 Meridian 23	0E		Acreas	ge 3242	
	Ecosite	042CY025NM SHALLOW SD-3			Photo Take	en Y	
W	vatershed	13060007110 COTTONWOOD- WALNUT					
С	bservers	NAVARRO/MCGEI	E	Obse	ervation Da	te 07/27/2	2004
County So	il Survey	NM666 CHAVES SO	OUTH	So	il Var/Taxa	ıd	
Soil N	Map Unit	Tg		Soil	Taxon Nan	ne TENC	EE
Texti	ure Class	NM666 GR-L			Soil Phas	TENC: UPTO	
Texture	Modifier	NM666 GRAVELLY LOAM	Z			· .	
	ved Avg Annual cipitation			Observed Avg Growing Season Precipitation		on	
	A Annual cipitation		10.42	NOAA Growing Season Precipitation		7.68	
NOAA Avg	g Annual cipitation	1/3/		NOAA Avg Growing Season Precipitation		- 1	10.24
	nces and mal Use:						
Part 2. Attı	ributes a	nd Indicators					
					ological Sit		S
Attribute	Indicator	TS	Extrem e	m Moderat e to Moderat e Moderat e		Slight to Moderat e	None to Slight
SH	Rills						X
Comments :					<u> </u>		
SH	Water Fl	ow Patterns				X	
Comments :							

SH	Pedestals and/or Terracettes			X	
Comments :					
SH	Bare Ground			X	
Comments :	50% rock with 20% rock cover	·.			
SH	Gullies			X	
Comments :					
S	Wind-scoured, Blowouts, and/or Deposition Areas			X	
Comments :					
Н	Litter Movement			X	
Comments :	Some displacement.				
SHB	Soil Surface Resistance to Erosion			X	
Comments :					
SHB	Soil Surface Loss or Degradation			X	
Comments :	Some pebbles and gravel have been some A-horizon loss.	migrated towards	the surface	and there l	nas
Н	Plant Community Composition and Distribution Relative to Infiltration and Runoff			X	
Comments :	Good infiltration, Grass cover is moisture along with pebbles an		oots are hole	ding the	
SHB	Compaction Layer			X	
Comments :					
В	Functional/Structural Groups			X	
Comments :	This is typical of an upland are	a with good grass	and shrub c	over.	
В	Plant Mortality/Decadence				X
Comments :					

НВ	Litter Amount				X	
Comments :	Litter amount falls well within	range.				
В	Annual Production			X		
Comments :	About 60% of potential and 50	0% of lon	g-term ave	erage.		
В	Invasive Plants			X		
Comments :	Creosote and Christmas cholla	scattered	d.			
В	Reproductive Capability of Perennial Plants					X
Comments :						
S	Physical/Chemical/Biologica l Crusts			X		
Comments :	Physical crusts especially in ir	nterspaces	S.			
В	Wildlife Habitat				X	
Comments :	Rolling hills desert grassland le.g., gravelly soils with creososites reflect current precipitation years). Draw bottoms within the	ote bush. On regime	Grass and fes (drought	forb producty over the	ction on sh past severa	allow
В	Wildlife Populations				X	
Comments :	No specific wildlife population are pronghorn antelope and a			• •		
В	Special Status Species Habitat					X
Comments :	None known to occur.					
В	Special Status Species Populations					X
Comments :	None known to occur.					
Part 3. Sun	nmary					
attributes be	r Summary - Each of the indica elow. An indicator is placed in Standard Attributes.					

Standard Attribute		Extrem e	Moderat e to Extreme	Moderat e	Slight to Moderat e	None to Slight
S	Soil	0	0	1	8	1
Н	Hydrologic	0	0	0	10	1
В	Biotic	0	0	2	7	4

B. Attribute Summary. In this table, the Extreme and Extreme to Moderate columns in the table above are merged for the *Does not Meet* column, Moderate becomes *May Need More Info*, and Slight to Moderate and None to Slight merge to form the *Meets* columns. Values from the table are summarized below. Space is provided for rationale of the determination. This space should most certainly be used when the determination by the ID team conflicts with the summarized values. Provide the sources of information that lead to the determination. X out the appropriate box for each attribute to denote final agreed upon determination by the ID team.

Attribute	Rationale	Does Not Meet	May Need More Info	Meet
Soil		0	1	9
Hydrologic		0	0	11
Biotic		0	2	11

Site Notes: Livestock are currently in this pasture but are utilizing the bottoms. There are natural drainage patterns occurring with sedimentation and erosion. From this upland area though the runoff is minimal.

RFOs	Upland	and Biotic Standa	rd Asse	ssment S	ummary `	Worksh	eet
		SITE 64079	-SOUT	H-F061			
Legal La	and Desc	NESW 18 0150S 024 Meridian 23	10E		Acreas	ge 2597	
	Ecosite	042CY025NM SHAI SD-3	LLOW		Photo Take	en Y	
W	atershed	13060007110 COTTONWOOD- WALNUT					
С	bservers	NAVARRO/MCGEF	Ξ	Obse	ervation Da	te 07/27/2	2004
County So	il Survey	NM666 CHAVES SO	OUTH	So	il Var/Taxa	ıd	
Soil N	Map Unit	Tg		Soil '	Taxon Nam	ne TENC	EE
Text	ure Class	NM666 GR-L			Soil Phas	TENCI UPTO	
Texture	Modifier	NM666 GRAVELLY LOAM	Z .				
	ved Avg Annual cipitation			Observed Avg Growing Season Precipitation		n	
	A Annual cipitation		10.42		AA Growin		7.68
NOAA Avg	g Annual cipitation	12.52		NOAA Avg Growing Season Precipitation		-	10.24
	nces and mal Use:					· .	
Part 2. Attı	ributes aı	nd Indicators					
					ological Sit		S
Attribute	Indicator	rs	Extrem e	Moderat Moderat Slight		Slight to Moderat e	None to Slight
S H	Rills						X
Comments			L		1		
S H	Water Fl	ow Patterns				X	
Comments :					',		

SH	Pedestals and/or Terracettes X
Comments :	
SH	Bare Ground X
Comments :	Now at 50%
SH	Gullies X
Comments :	
S	Wind-scoured, Blowouts, and/or Deposition Areas
Comments :	
Н	Litter Movement X
Comments :	
SHB	Soil Surface Resistance to Erosion X
Comments :	
SHB	Soil Surface Loss or Degradation
Comments :	
Н	Plant Community Composition and Distribution Relative to Infiltration and Runoff
Comments :	
SHB	Compaction Layer X
Comments :	
В	Functional/Structural Groups X
Comments :	Absence of gramas and dropseed.
В	Plant Mortality/Decadence X
Comments :	Creosote appears to be treated. Christmas cholla is growing up from most of treated decadent creosote plants.

НВ	Litter Amount				X	
Comments :	Now 10-20%					
В	Annual Production			X		
Comments :	Production is now estimated a	t 40% of	long-term	average.		
В	Invasive Plants			X		
Comments :	Snakeweed common and creos	sote scatt	ered.			
В	Reproductive Capability of Perennial Plants				X	
Comments :						
S	Physical/Chemical/Biologica l Crusts				X	
Comments :	Physical crusts evident.					
В	Wildlife Habitat				X	
Comments :	Rolling hills desert grassland le.g., gravelly soils with creoso sites reflect current precipitation years). Draw bottoms within the Note: This pasture was chemically contained to the contained	te bush. On regime	Grass and fes (droughte are the m	Forb productions ore productions or the contraction of the contraction	etion on sh past severative sites.	allow al
В	Wildlife Populations				X	F 112 11
Comments :	No specific wildlife population pronghorn antelope and a varie					
В	Special Status Species Habitat					X
Comments :	None known to occur.					
В	Special Status Species Populations					X
Comments :	None known to occur.					
Part 3. Sun	nmary					
	Summary - Each of the indicated with Summary Summary - Each of the indicator is placed in a					


each of the	Standard Attributes.					
Standard Attribute		Extrem e	Moderat e to Extreme	Moderat e	Slight to Moderat	None to Slight
S	Soil	0	0	0	8	2
Н	Hydrologic	0	0	0	9	2
В	Biotic	0	0	3	7	3

B. Attribute Summary. In this table, the Extreme and Extreme to Moderate columns in the table above are merged for the *Does not Meet* column, Moderate becomes *May Need More Info*, and Slight to Moderate and None to Slight merge to form the *Meets* columns. Values from the table are summarized below. Space is provided for rationale of the determination. This space should most certainly be used when the determination by the ID team conflicts with the summarized values. Provide the sources of information that lead to the determination. X out the appropriate box for each attribute to denote final agreed upon determination by the ID team.

Attribute	Rationale	Does Not Meet	May Need More Info	Meet
Soil		0	0	10
Hydrologic		0	0	11
Biotic		0	3	10

Site Notes: This site is upland from a dirt tank where mesquite is dominate. A pipeline intersects two of the transects. Photos were taken.

### **Ground Cover Trends**



■ SROCK

Forb
BGROUND

Shrub
LROCK
LITTER
Grass

Λ.	<i>''D'</i>			<sup>7</sup> /O,
	11/1/1982	10/28/1986	10/25/1991	11/2/2001
BGROUND	23.00	75.00	50.00	55.00
Forb	3.00	0.00	3.00	0.00
Grass	18.00	15.00	16.00	7.00
LITTER	12.00	2.00	8.00	3.00
LROCK	3.00	0.00	2.00	0.00
Shrub	7.00	7.00	0.00	6.00
SROCK	32.00	0.00	2.00	29.00

	11/1/1982	10/28/1986	10/25/1991	11/2/2001
Total	98.00	99.00	81.00	100.00

### **Report Parameters**

SITE NAME LIKE 64079-NORTH-F060

ON/AFTER 10/01/1982 ON/BEFORE 09/30/2002

# **Functional / Structural Groups**

#### **Report Parameters**

SITE NAME LIKE 64079-NORTH-F060


ON/AFTER 10/01/1982 ON/BEFORE 09/30/2002

MIN LBS TO GRAPH 1


SELECTED ECOSITE 042CY025NM

Group	Plant Type	Species	Low Wt Allowed	High Wt Allowed	Minimum	Maximum	Average	STDEV
1	Grass	BOER4	105	157	12.00	201.00	76.43	63.63
2	Grass	BOCU	78	105	0.00	4.00	0.67	1.49
3	Grass	BOGR2	78	105	0.00	9.00	2.67	3.54
4	Grass	MUPO2	26	52	0.00	26.00	6.50	11.26
6	Grass	SPCR	26	52	0.00	29.00	6.67	10.19
7	Grass	TRMU	15	26	0.00	110.00	25.71	36.50
7	Grass	TRPI2	15	26	0.00	37.00	14.83	12.24
10	Grass	ERPU8	5	15	1.00	55.00	13.00	17.90
11	Grass	ARIST	15	26	2.00	107.00	35.71	31.54
11	Grass	HIMU2	15	26	0.00	37.00	7.67	13.20
11	Grass	MUAR2	15	26	0.00	46.00	17.67	14.62
11	Grass	SCBR2	15	26	0.00	49.00	12.71	15.37
14	Grass	PAHA	5	15	0.00	4.00	0.67	1.49
17	Forb	SPHAE	5	15	0.00	0.00	0.00	0.00
20	Forb	CROTO	10	26	0.00	16.00	6.43	5.39
20	Forb	CRPO5	10	26	0.00	3.00	0.60	1.20
20	Forb	DYAC	10	26	0.00	17.00	2.83	6.34
21	Forb	AAFF	5	15	0.00	22.00	8.75	9.31
21	Forb	CARLI	5	15	0.00	1.00	0.17	0.37
21	Forb	DYPE	5	15	0.00	56.00	17.80	22.70
22	Forb	COCA2	5	15	0.00	3.00	0.50	1.12
22	Forb	ERTE13	5	15	0.00	3.00	0.60	1.20
22	Forb	HOGL2	5	15	0.00	2.00	0.67	0.75
22	Forb	MELE2	5	15	0.00	7.00	2.00	2.76
22	Forb	PENA	5	15	0.00	1.00	0.50	0.50
25	Shrub	LADI2	5	15	0.00	66.00	27.00	20.78

Group	Plant Type	Species	Low Wt Allowed	High Wt Allowed	Minimum	Maximum	Average	STDEV
26	Shrub	KRPA	5	15	0.00	3.00	1.20	1.47
27	Shrub	COER5	5	15	0.00	16.00	4.75	6.61
32	Shrub	OPLE	5	15	0.00	6.00	2.00	2.83
32	Shrub	OPUNT	5	15	0.00	1.00	0.17	0.37
34	Shrub	GUSA2	5	15	0.00	106.00	53.83	37.79
36	Shrub	ACACI	5	15	0.00	4.00	2.00	2.00
36	Tree	ACGR	5	15	0.00	2.00	0.40	0.80
36	Shrub	DAFO	5	15	0.00	28.00	5.57	9.41
36	Shrub	DYPE3	5	15	0.00	102.00	46.67	42.09
36	Shrub	MESC	5	15	0.00	4.00	1.33	1.89
36	Shrub	ZIPU	5	15	0.00	21.00	3.50	7.83



### **Production Lbs/Acre Trends**




	11/1/1982	11/2/1983	1/18/1985	11/22/1985	10/28/1986	10/25/1991	11/2/2001
Forb	31.00	66.00	64.00	11.00	31.00	159.00	0.00
Grass	139.00	79.00	149.00	163.00	447.00	422.00	98.00
Shrub	195.00	121.00	86.00	78.00	199.00	79.00	41.00
Tree	0.00	2.00	0.00	0.00	0.00	0.00	10.00
Total	365.00	268.00	299.00	252.00	677.00	660.00	149.00

### **Report Parameters**

SITE NAME LIKE 64079-NORTH-F060

ON/AFTER 10/01/1982 ON/BEFORE 09/30/2002

# **Ground Cover Trends**



	11/1/1982	10/28/1986	10/25/1991	11/2/2001
BGROUND	66.00	73.00	62.00	71.00
Grass	13.00	16.00	14.00	18.00
LITTER	12.00	4.00	9.00	3.00
Shrub	8.00	7.00	0.00	2.00
SROCK	0.00	0.00	15.00	6.00
Total	99.00	100.00	100.00	100.00

### **Report Parameters**

SITE NAME LIKE 64079-SOUTH-F061

ON/AFTER 10/01/1982 ON/BEFORE 09/30/2002

# **Functional / Structural Groups**

#### **Report Parameters**

SITE NAME LIKE 64079-SOUTH-F061


ON/AFTER 10/01/1982 ON/BEFORE 09/30/2002

MIN LBS TO GRAPH 1


SELECTED ECOSITE 042CY025NM

Group	Plant Type	Species	Low Wt Allowed	High Wt Allowed	Minimum	Maximum	Average	STDEV
1	Grass	BOER4	105	157	13.00	233.00	86.67	76.20
3	Grass	BOGR2	78	105	0.00	9.00	2.83	4.02
3	Grass	BOHI2	78	105	0.00	9.00	2.25	3.90
6	Grass	SPCR	26	52	0.00	41.00	10.00	14.41
7	Grass	TRMU	15	26	0.00	38.00	13.83	13.50
7	Grass	TRPI2	15	26	0.00	34.00	13.67	10.31
8	Grass	MUAR	5	15	0.00	8.00	3.33	2.98
10	Grass	ERPU8	5	15	4.00	68.00	22.83	21.57
11	Grass	ARIST	15	26	0.00	89.00	28.33	28.83
11	Grass	HIMU2	15	26	0.00	32.00	15.40	11.89
11	Grass	MUAR2	15	26	0.00	31.00	10.25	12.17
11	Grass	SCBR2	15	26	4.00	66.00	27.80	21.34
12	Grass	AAGG	0	5	0.00	1.00	0.50	0.50
18	Forb	LESQU	5	15	0.00	1.00	0.50	0.50
20	Forb	CROTO	10	26	0.00	10.00	3.00	3.79
20	Forb	CRPO5	10	26	0.00	5.00	1.25	2.17
20	Forb	DYAC	10	26	0.00	0.00	0.00	0.00
21	Forb	AAFF	5	15	0.00	2.00	1.00	0.82
21	Forb	COLDE	5	15	0.00	0.00	0.00	0.00
22	Forb	ERTE13	5	15	0.00	5.00	1.80	2.23
22	Forb	HOGL2	5	15	0.00	1.00	0.20	0.40
22	Forb	MELE2	5	15	0.00	8.00	2.20	3.12
22	Forb	PENA	5	15	0.00	1.00	0.25	0.43
25	Shrub	LADI2	5	15	3.00	88.00	45.00	32.84
32	Shrub	OPIM	5	15	0.00	4.00	1.33	1.89
34	Shrub	GUSA2	5	15	11.00	242.00	113.40	74.86

Group	Plant Type	Species	Low Wt Allowed	High Wt Allowed	Minimum	Maximum	Average	STDEV
36	Tree	ACGR	5	15	0.00	2.00	0.50	0.76
36	Shrub	DAFO	5	15	0.00	17.00	2.83	6.34



### **Production Lbs/Acre Trends**



	11/1/1982	11/2/1983	11/22/1985	10/28/1986	10/25/1991	11/2/2001
Forb	10.00	14.00	1.00	14.00	157.00	0.00
Grass	89.00	79.00	151.00	413.00	409.00	216.00
Shrub	211.00	325.00	97.00	143.00	90.00	36.00
Tree	1.00	0.00	0.00	0.00	2.00	0.00
Total	311.00	418.00	249.00	570.00	658.00	252.00

### **Report Parameters**

SITE NAME LIKE 64079-SOUTH-F061

ON/AFTER 10/01/1982 ON/BEFORE 09/30/2002



# Rangeland Health Assessment Ecological Sites



No tenently is more by the Ennes inflict of Mercapolitances with the Gardinery pointed by a strongly of oscial discounts from the purposes not time shall by ELM. Spott for Normal tenently on the purposes not time shall by ELM. Spott for Normal tenently of the Normal N

Allotment 64093

64079


T14.R23E

Study Plots

Private

Study Locations

Public



**Ecological Sites** 

Allotment Boundary



# Rangeland Health Assessment **Soil Mapping Units**



Allotment <del>6409</del>3


64079

T14.R23E

Public

Private

Study Locations



Soil Mapping Units

Allotment Boundary

GIS Intern on July 25, 2003.

No tenently is more by the Ennes inflict of Mercapolitances with the Gardinery pointed by a strongly of oscial discounts from the purposes not time shall by ELM. Spott for Normal tenently on the purposes not time shall by ELM. Spott for Normal tenently of the Normal Vap According European This is format for many boundard with an incoming tenent.